

Microblading Pre-Procedure Advice

PLEASE READ THE FOLLOWING ADVICE CAREFULLY AND SIGN AT THE END

- Microblading procedure generally requires 2 sessions for best results. Client will return for a touch-up in 4-6 weeks after initial procedure. Those with oily skin may require an additional touch-up. Please be aware that color intensity will be significantly darker and sharper immediately after the initial procedure and will reduce by 30-40%.
- Delicate or sensitive skin may be red following the procedure.
- Please do not drink alcohol 24 hours prior to the treatment.
- Any brow waxing or threading should be done at least 72 hours before procedure.
- AHA products and retinoids should be avoided no less than 2 weeks prior to procedure.
- Exfoliating treatments such as microdermabrasion should not be performed within 2 weeks prior to procedure.
- Chemical and laser peels should be avoided no less than 6 weeks prior to procedure.
- Please wait 2 months if you have had Botox treatment before microblading.
- Do not drink coffee or anything with caffeine the day of procedure. Do not take Aspirin, Vit. E, or ibuprofen 48 hours prior to procedure.

TOPICAL ANESTHETIC ADVICE

- This is designed to be a gentle procedure with minimal discomfort. The degree of discomfort will depend on your pain threshold. We do apply a topical 5% lidocaine after an initial outline of your brows. Once absorbed you should be comfortable with only a scratching feeling.
- You may purchase a topical numbing from your local pharmacy and apply the cream 30 min. prior to procedure if you're concerned about pain. This may give some relief.
- Some experience more pain during their menstrual cycle.
- Allergic reactions are always a possibility. We will discuss any allergies you have during the consult. Reactions may occur after the procedure and may show through rash, blistering, increasing redness or any other symptoms associated with allergic reaction. If this occurs please consult your physician.

CONTRAINDICATIONS FOR MICROBLADING

- Liver disease
- Compromised skin near the brow area
- Hormone therapies can affect pigmentation and/or cause sensitivity.
- Pregnancy/Nursing
- Chemotherapy/Radiation
- Any medical disease that causes slow healing or high risk infection such as diabetes, auto-immune disease, Graves' disease etc. A clearance from your doctor is recommended.
- Previous tattoos.....consult required.

I HAVE READ THE INFORMATION PROVIDED AND FULLY UNDERSTAND ANY RISKS. I AGREE TO FOLLOW THE PRE AND POST PROCEDURE ADVICE.

Client Signature _____ Date _____